

## WARRANTY

Products are guaranteed to conform to the quality and content indicated on each vial and external labels during their shelf life. BIOTOOLS obligation and purchaser's rights under this warranty are limited to the replacement by BIOTOOLS of any product that is shown defective in fabrication, and that must be returned to BIOTOOLS, freight prepaid, or at BIOTOOLS' option, replacement of the purchasing price. Any complaint on damaged goods during transport must be directed to the handling or transport agent.

Product for Research Use Only. This product must be used by qualified professionals only. It is the responsibility of the user to ascertain that a given product is adequate for a given application. Any product not fulfilling the specifications included in the product sheet will be replaced. This warranty limits our responsibility to the replacement of the product. No other warranties, of any kind, express or implied, including, without limitation, implicit warranties of commercialisation ability or adequacy for a given purpose, are provided by BIOTOOLS. BIOTOOLS will not be held responsible for any direct, indirect, consequential or incidental damage resulting of the use, misuses, results of the use or inability to use any product.

### Produced by:

BIOTOOLS, Biotechnological & Medical Laboratories, S.A. have been evaluated and certified to accomplish ISO 9001:2008 requirements for the following activities: Research and development of biotechnology products and manufacture of biotechnology and in vitro products. Valle de Tobalina – 52 – Nave 39, 28021 Madrid – Spain

© 2008 BIOTOOLS, Biotechnological & Medical Laboratories, S.A. All rights reserved



BIOTOOLS B&M Labs, S.A.  
Valle de Tobalina - 52 - Nave 39  
28021 Madrid  
Spain

Tel. (34) 91 710 00 74  
Fax (34) 91 505 31 18  
E-mail: [info@biotools.eu](mailto:info@biotools.eu)  
[www.biotools.eu](http://www.biotools.eu)



# BIOTOOLS

BIOTOOLS B & M LABS. S.A.

## BIOTOOLS *Pfu* DNA POLYMERASE (1 U/μl)

REF.	FORMAT	CONTENT
10.501	100 U	BIOTOOLS <i>Pfu</i> DNA Polymerase (1 U/μl) 10X Standard Reaction Buffer with MgCl <sub>2</sub>
10.502	250 U	BIOTOOLS <i>Pfu</i> DNA Polymerase (1 U/μl) 10X Standard Reaction Buffer with MgCl <sub>2</sub>
10.511	100 U	BIOTOOLS <i>Pfu</i> DNA Polymerase (1 U/μl) 10X Reaction Buffer MgCl <sub>2</sub> FREE
10.512	250 U	BIOTOOLS <i>Pfu</i> DNA Polymerase (1 U/μl) 10X Reaction Buffer MgCl <sub>2</sub> FREE
10.500BW	BULK	BIOTOOLS <i>Pfu</i> DNA Polymerase (1 U/μl)
10.500B	BULK	BIOTOOLS <i>Pfu</i> DNA Polymerase (1 U/μl) 10X Standard Reaction Buffer with MgCl <sub>2</sub>
10.510B	BULK	BIOTOOLS <i>Pfu</i> DNA Polymerase (1 U/μl) 10X Reaction Buffer MgCl <sub>2</sub> FREE

Store at -20°C

**Notice to users:** Some of the applications which may be performed with this product are covered by applicable patents in certain countries. The purchase of this product does not include or provide a license to perform patented applications. Users may be required to obtain a license depending on the country and/or application.

Ed 12 – Sept. 2021

## 1. GENERAL CONSIDERATIONS

BIOTOOLS *Pfu* DNA Polymerase is a highly thermostable polymerase with proof-reading activity. In addition to 5'-3' DNA polymerase activity, it also possesses 3'-5' exonuclease (proofreading) activity. It is a recombinant protein isolated from *Pyrococcus furiosus* expressed in *E. coli*.

BIOTOOLS *Pfu* DNA polymerase is recommended for use in PCR and primer extension reactions that require high fidelity. *Pfu* DNA polymerase has an error rate 10-fold lower than non proof-reading DNA polymerases.

The enzyme is free of unspecific endonuclease activity, as well as nicking activities. It does not exhibit nucleotidyl terminal transferase activity so its amplification products can be directly used for cloning in blunt-ended vectors.

The enzyme is supplied at a **concentration of 1 U/μl** in a storage buffer. This concentration allows accurate pipetting of small amounts of the DNA polymerase, so that further dilutions are not necessary.

### Product Applications:

- High-fidelity PCR
- Long and complex amplifications
- Gene cloning
- Mutation analysis
- Standard PCR
- In situ PCR

## 2. ENZYME FEATURES

Concentration: .....	1 U/μl
Optimal activity:	
Enzyme concentration.....	20-50 mU/μl
pH .....	8-9
Elongation temperature.....	72-75 °C
MgCl <sub>2</sub> concentration .....	2 mM
Size of PCR products: .....	Up to 5 Kb
PCR cloning: .....	Blunt ends
Endonuclease activity: .....	No
Reverse transcriptase activity: .....	No
5'→3' exonuclease activity: .....	No
3'→5' exonuclease activity: .....	Yes
Nicking activity: .....	No

This enzyme is not recommended for certain experiments dealing with amplification of sequences homologous to those found in *E. coli* or amplifications with very low annealing temperatures (e.g. RAPDs, Random Amplified Polymorphic DNAs)

## 3. STORAGE CONDITIONS

Store package components at -20°C in a constant temperature freezer. If stored under the recommended conditions, the product will maintain performance through the indicated date on the label.

## 4. PRODUCT SPECIFICATIONS

**Unit Definition-** One unit is defined as the amount of enzyme which incorporates 10 nanomoles of dNTPs into acid-insoluble DNA within 30 minutes at 72 °C.

**Storage buffer-** 20 mM Tris-HCl (pH 8.0); 50 mM KCl; 0.25 % NP 40; 0.25 % Tween 20; 40 % glycerol (v/v).

**10X Reaction Buffer-** 750 mM Tris HCl (pH 9.0); 500 mM KCl; 200 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>. The **10X STANDARD REACTION BUFFER with MgCl<sub>2</sub>** also includes 20 mM MgCl<sub>2</sub> in its composition.

## 5. GENERAL ASPECTS OF REACTION COMPONENTS

### Enzyme Concentration

Biotools *Pfu* DNA Polymerase is suitable for standard and specialized PCR applications. As an initial guide we recommend employing the following enzyme units per reaction.

Final reaction volume	Recommended enzyme units
100 μl	2.0-2.5 Units
50 μl	1.0-1.25 Units
25 μl	0.5-0.75 Units

The addition of higher quantities of enzyme generally does not produce significant yield increase. Only for certain applications (e.g. PRIND or Primed In Situ Synthesis) or when working on long DNA fragment amplifications (longer than 2 Kb from genomic DNA) it might be necessary to increase the concentration.

### DNA Template

The quality and quantity of the DNA template affects both the sensitivity and efficiency of the amplification. High amounts of DNA usually increase the amplification of nonspecific PCR products. The PCR is inhibited by various compounds e.g. ionic detergents, phenol, gel loading dyes, etc. If the DNA template contains traces of inhibitors, reduce the amount of the template included in the amplification reaction, or repurify the template by ethanol precipitation and several washing steps.

### dNTPs Concentration

The concentration of each dNTP in amplification reactions should be 50-500 μM, being 200 μM the most commonly used concentration. The concentration of dNTPs may be decreased (e.g. when unspecific amplification occurs), increased (e.g. for long amplifications), or even unbalanced in favour of any of the particular dNTPs (e.g. in vitro mutagenesis experiments).

The dNTPs behave as potent Mg<sup>2+</sup> chelating agents reducing therefore the availability of free Mg<sup>2+</sup> for polymerase activity. Thus an increase in dNTPs should be accompanied by an increase in MgCl<sub>2</sub> concentration.

Biotools *Pfu* DNA Polymerase is suitable for its use with modified dNTPs (e.g. radioactively or fluorescein labelled) as substrate. It can also be used with dUTP and other analogues.

### Reaction Buffer

The provided buffer has been specially formulated to facilitate the amplification of any PCR products. It creates the appropriate stringent conditions for primer-annealing over a wide range of temperatures. Moreover, the **Standard Reaction Buffer** includes Mg<sup>2+</sup> at the optimal concentration for most experiments (final concentration: 2 mM) which facilitates the use of the enzyme.

## 7. GUIDE TO AMPLIFICATION PROGRAM

### MgCl<sub>2</sub> Concentration

The optimal MgCl<sub>2</sub> concentration may vary depending on the primer and template that are used and must be determined by experimentation. In most cases, a final concentration of MgCl<sub>2</sub> at 2 mM in the reaction mix works well.

High concentrations of MgCl<sub>2</sub> may promote low enzyme fidelity and non specific amplification products; whereas low concentrations should reduce the yield of the desired amplification products. If the samples contain any chelating metal agents such as EDTA, the concentration of MgCl<sub>2</sub> should be increased accordingly.

### Primer Design

The proofreading activity of *Pfu* DNA Polymerase may degrade the amplification primers resulting in nonspecific amplification and reduced product yield. To overcome primer degradation, longer primers (20-35 bases) with maximized G+C content can be used. Primers can also be protected by introducing a single phosphorothioate bound at their 3' end.

To avoid primer-dimer and hairpin formation the primers should not be self-complementary or complementary to any other primer present in the reaction. The annealing temperature of the primers should be similar (< 5°C variation).

The 5' end of a primer may contain mismatches between the primer and template, whereas this is not recommended at the 3' end.

### PCR Additives

In complex amplifications the presence of DMSO, betaine, formamide or any other PCR additives might be necessary. The provided enzyme and buffer are compatible with most PCR additives. When calculating the annealing temperature for the PCR cycling program, it is important to take into account that certain additives may decrease the melting temperature of the primers.

## 6. STANDARD PROTOCOL

Optimal conditions must be determined for each individual experimental system.

**Proceed to the Reagent Preparation Area in a laminar flow cabinet.**

1. Thaw and keep reagents on ice. After complete thawing, mix the reagents well and spin down in a bench-top centrifuge.
2. Prepare a master mix according to Table 1. For each experiment include at least one negative control (without template). To ensure sufficient volume include additional reactions in the calculations.

**Note 1: Adding the *Pfu* DNA polymerase at the end is recommended to avoid oligonucleotide degradation because of its 3'-5' exonuclease activity, maximum is added in the absence of dNTPs.**

TABLE 1. Master Mix Preparation

COMPONENTS	Final Concentration	50 µl rxn	20 µl rxn
<b>Master Mix</b>			
10X REACTION BUFFER	1X	5 µl	2 µl
50 mM MgCl <sub>2</sub> *	1.5-4 mM	1.5-4 µl	0.6-1.6 µl
dNTP Mix 10 mM each	200 µM of each	1 µl	0.4 µl
Primers	variable	variable	variable
Nuclease-free water	-	To 50 µl	To 20 µl
<i>Pfu</i> DNA Polymerase (1U/µl)**	20-50 mU/µl	1.0-2.5 µl	0.4-1.0 µl
<b>Template DNA</b>	<b>Variable</b>	<b>Variable</b>	<b>Variable</b>

\*not necessary for 10X Standard Reaction Buffer, it includes MgCl<sub>2</sub>

\*\*See Note 1

3. Mix the master mix thoroughly and keep on ice. Distribute the appropriate volume into each vial.

**Proceed to DNA Purification Area separate from other sources of DNA.**

4. Add the template DNA to each reaction vial. Close the vials and mix gently. For thermal cycler without heated lid overlay a mineral oil layer.

**Proceed to the Amplification Area**

5. Program the thermal cycler according to Table 2. Place the vials in the thermal cycler and perform the selected PCR program.

TABLE 2. Standard Amplification Program

Cycle Step	Nº Cycles	Temperature	Time
Initial Denaturation	1	94°C	3-10 min*
Denaturation	25-35**	94°C	5-30 sec
Annealing		T <sub>m</sub> -5°C	30-60 sec
Extension		72°C	2 min/1 kb
Final Extension	1	72°C	5-15 min
Cooling	∞	4°C	∞

\*Depending on the template (see Point 7).

\*\*Optimize time; temperature; and number of cycles (see Point 7).

**Initial Denaturation Step-** Incomplete denaturation of the PCR reaction results in an inefficient first amplification cycle and low amplification yield. However, the denaturation must be kept as short as possible in order to avoid inactivation of the enzyme. For most samples 94°C for 3-5 mins should be satisfactory. Templates rich in G+C content often require longer initial denaturation and the length of this step can be extended (up to 10 minutes).

**Denaturation Step-** The PCR product synthesized in the amplification cycling is shorter than the template DNA and therefore needs a short denaturation step; 5-30 seconds of denaturation at 94°C should be sufficient.

**Primer Annealing Step-** To find the optimal annealing temperature, you can use a temperature gradient. Start using an annealing temperature 5 °C below the T<sub>m</sub> of the primers. If the primers have a high T<sub>m</sub> a two-step cycling is recommended.

**Extension Step-** The annealed primers must be extended at 70-75°C. The extension time depends on the size of the expected product. Due to the proofreading activity of the *Pfu* DNA Polymerase, it requires more time for extension as compared to other polymerases. An extension time of 2 min for every 1 kb to be amplified should be used.

**Number of PCR Cycles-** Cycling program usually consists on 25-35 cycles. This parameter depends on the amount of starting material and the expected yield. In certain assays, increasing the number of cycles leads to an increase in nonspecific products and consequently to a decrease in the yield of specific product. You should experimentally determine the optimal number of cycles for each experiment.

**Final Extension Step-** After the last PCR cycle the sample should be incubated at 72°C for 5-15 min. The *Pfu* DNA polymerase does not add extra adenine nucleotides to the 3' ends of the PCR products.

## 8. TROUBLESHOOTING

Problem	Cause	Recommendation
Low yield or no amplification product	Missing reagent or pipetting error	Check concentration and storage conditions of dNTPs, primers, etc. Repeat the PCR including all reagents.
	DNA template problems	Check the concentration and quality of starting material. Repeat the PCR with a new dilution of template or with a new DNA purification. If the template is difficult e.g. rich in G+C sequences we recommend adding DMSO to the master mix.
	Problems with primers	Revise the primers design and the primers storage condition. Check primer degradation on a denaturing polyacrylamide gel. Avoid any design prone to the formation of primer dimers. Repeat PCR with different primer concentration from 0.1-0.5 µM in 0.1 increments. It is critical to add the <i>Pfu</i> DNA Polymerase to the amplification reaction last, particularly following the addition of dNTP (see Note 1).
	Enzyme concentration too low	Increase enzyme concentration in 0.2 U increments.
	MgCl <sub>2</sub> concentration	Optimise MgCl <sub>2</sub> concentration of the PCR if necessary (1.5-4 mM)
Nonspecific amplification products or background smear	Incorrect PCR cycling conditions	Check the following parameters of the PCR program (see point 7): <b>Denaturation</b> - Increase time and temperature of initial denaturation. <b>Annealing</b> - Optimise the annealing temperature and time. <b>Extension time</b> - Increase extension time by increments of 30 sec. <b>Number of cycles</b> - Perform additional cycles by increments of 5 cycles.
	Problems with primers	Check primer degradation on a denaturing polyacrylamide gel. Design alternative primers. Both primers should be present at the same concentration (0.1-0.5 µM). Decrease primer concentration by increments of 0.1 µM.
	Excess of DNA template	Use dilutions of your template.
	Enzyme concentration too high	Optimise polymerase concentration of the PCR if necessary
	MgCl <sub>2</sub> concentration	Optimise MgCl <sub>2</sub> concentration of the PCR if necessary (1.5-4 mM)
PCR products in negative control	Incorrect PCR cycling	Increase the annealing temperature in increments of 1°C and/or reduce the time of this step. Reduce the number of cycles.
	Carryover contamination	Exchange all reagents.

## 9. ORDERING INFORMATION

Components	References						
	10.501	10.502	10.511	10.512	10.500BW	10.500B	10.510B
<b>Biotoools <i>Pfu</i> DNA Polymerase (1 U/µl)</b>	100 U	250 U	100 U	250 U	On demand	On demand	On demand
<b>10X Standard Reaction Buffer with MgCl<sub>2</sub></b>	1.8ml	1.8ml				On demand	
<b>10X Reaction Buffer MgCl<sub>2</sub> FREE</b>			1.8ml	1.8ml			On demand
<b>50 mM MgCl<sub>2</sub> Solution</b>			1.8ml	1.8ml			

